



Leafy chenopods: a nutritional powerhouse

Niraj Kumar Prajapati^a, Suraj Pasawan^a, Maya Ram^a and Sanjay Kumar^a

Summary

Leafy chenopods, a group of nutrient-dense, underutilized plant species, have emerged as a promising nutritional powerhouse with immense potential to combat micronutrient malnutrition globally. These resilient, drought-tolerant plants, including quinoa, amaranth, and huauzontle, have been cultivated for centuries by indigenous communities in the Americas and are now gaining recognition for their exceptional nutritional profile. Leafy chenopods are rich sources of protein, dietary fibre, and essential amino acids, making them valuable additions to plant-based diets. Moreover, they are abundant in micronutrients such as iron, zinc, calcium, and vitamins A, C, and E, addressing common deficiencies in many populations. Their high antioxidant content, derived from compounds like betalains and polyphenols, may confer protective effects against chronic diseases like cancer, cardiovascular disorders, and diabetes. Additionally, their anti-inflammatory and antimicrobial properties hold promise for therapeutic applications. These resilient crops can thrive in marginal lands and harsh climatic conditions, making them well-suited for cultivation in areas affected by climate change and water scarcity. Their tolerance to various biotic and abiotic stresses further enhances their potential as sustainable and climate-smart crops. Leveraging these underutilized crops could be a significant step towards achieving food and nutrition security in the face of climate change and population growth.

JAE 2024, Vol 18

Received: 28 February 2024

Accepted: 25 April 2024

Published: 29 May 2024

<https://doi.org/10.58628/JAE-2418-104>

Associated Editor: Dr. SM Haldhar

Copyright © 2024 The Author(s). Published by Society for Agriculture and Arid Ecology Research (SAAER). This is an Open Access article under the Creative Commons Attribution License 4.0 (CC BY-NC-SA).



Keywords: Bathua, fat hen, pig-weed, leafy chenopod, nutritional, health benefits

Introduction

The plant family Amaranthaceae, also known as the amaranth family, encompasses a diverse group of flowering plants widely distributed across the globe. Within this family, the chenopod genus (*Chenopodium*) stands out as a treasure trove of nutritious, leafy greens that various cultures have long consumed. These unassuming plants, often overlooked and underappreciated, are emerging as nutritional powerhouses packed with an array of essential vitamins, minerals, and beneficial compounds. This review delves into the remarkable nutritional profile of leafy chenopods, exploring their potential to address global nutritional challenges and promote overall well-being.

The chenopod genus comprises over 150 species, many of which are herbaceous annuals or perennials¹. These plants are commonly known by various names, including lambs' quarters, goosefoot, fat-hen, and pigweed. While some species are considered weeds, others have been cultivated for centuries as nutrient-dense leafy greens or pseudocereals².

Nutritional Composition

Leafy chenopods are renowned for their exceptional nutritional value, boasting an impressive array of essential nutrients. Their nutrient profile is comparable

to, and in some cases, surpasses that of many conventionally consumed leafy greens.

Macronutrients

Proteins: Leafy chenopods are rich in high-quality proteins, containing a well-balanced amino acid profile³. The protein content can range from 2.5% to 5% of the fresh weight, making them a valuable source of plant-based protein⁴.

Carbohydrates: These leafy greens are low in carbohydrates, with a predominance of dietary fibre, which contributes to their potential benefits for digestive health and glycaemic control⁵.

Lipids: While not exceptionally high in total fat content, leafy chenopods contain a favourable fatty acid profile, with a high proportion of essential polyunsaturated fatty acids, such as linoleic and linolenic acids⁶.

Micronutrients

Vitamins: Leafy chenopods are exceptional sources of various vitamins, including vitamin A (in the form of carotenoids like lutein and zeaxanthin), vitamin C, vitamin E, and several B vitamins, such as folate, riboflavin, and niacin^{5,7}.

Minerals: These nutrient-dense greens are rich in essential minerals, including calcium, iron, magnesium, phosphorus, potassium, and zinc^{5,7}. The bioavailability of these minerals is enhanced by the presence of organic acids and chelating compounds in the leaves⁸.

^aDepartment of Horticulture, School of Agricultural Sciences and Technology, Babasaheb Bhimrao Ambedkar University (A Central University), Lucknow (U.P.) – 226025, India

[#]Corresponding author NK Prajapati, E-mail: nkp.ofcl@gmail.com

Table 1. Nutritional Composition of Selected Leafy Chenopods (per 100g edible portion)⁹

Nutrient	Quinoa Leaves	Amaranth Leaves	Huauzontle Leaves
Protein (g)	4.4	4.8	5.2
Dietary Fiber (g)	5.2	6.7	7.1
Iron (mg)	6.5	7.6	8.2
Zinc (mg)	0.9	1.2	1.5
Vitamin A (µg RAE)	510	688	742
Vitamin C (mg)	48.1	61.5	75.2

Phytochemicals: Leafy chenopods are excellent sources of various phytochemicals, including flavonoids, phenolic acids, and betalains, which contribute to their antioxidant and anti-inflammatory properties^{5,6}.

Betalains, a class of pigments responsible for the vibrant colours of some chenopod species, have been associated with various health benefits, such as antioxidant, anti-inflammatory, and potential anti-cancer effects^{10,11}.

Antioxidant Properties: The abundance of phytochemicals, vitamins, and minerals in leafy chenopods contributes to their potent antioxidant capacity^{5,6}. Antioxidants play a crucial role in neutralising free radicals and mitigating oxidative stress, which has been implicated in various chronic diseases, such as cardiovascular disorders, neurodegenerative diseases, and certain cancers¹².

Table 2. Antioxidant activity and total phenolic content of leafy chenopods¹³

Plant Species	DPPH Radical Scavenging Activity (%)	Total Phenolic Content (mg GAE/g dry weight)
Quinoa Leaves	82.1 ± 2.3	12.6 ± 0.4
Amaranth Leaves	88.5 ± 1.8	17.3 ± 0.7
Huauzontle Leaves	91.2 ± 2.1	21.5 ± 1.1

Anti-inflammatory Effects: Several compounds found in leafy chenopods, including flavonoids, phenolic acids, and betalains, have been shown to possess anti-inflammatory properties^{5,6}. Chronic inflammation is a contributing factor to numerous health conditions, such as rheumatoid arthritis, inflammatory bowel diseases, and certain types of cancer¹⁴. Incorporating leafy chenopods into the diet may help modulate inflammation and mitigate associated health risks.

Cardiovascular Benefits: The favourable nutrient profile of leafy chenopods, including their high fibre content, unsaturated fatty acids, and antioxidant compounds, may contribute to improved cardiovascular health^{5,7}. These greens have been associated with the potential to lower blood pressure, improve lipid profiles, and reduce the risk of cardiovascular diseases⁴.

Digestive Health: The high fibre content and prebiotic properties of leafy chenopods can promote digestive health by facilitating regular bowel movements,

preventing constipation, and supporting a healthy gut microbiome^{5,15}. Additionally, the presence of beneficial compounds like betalains may exert protective effects against gastrointestinal disorders¹.

Diabetes Management: Due to their low glycaemic index and high fibre content, leafy chenopods may be beneficial for individuals with diabetes or those at risk of developing the condition^{4,5}. The fibre and compounds present in these greens can help regulate blood sugar levels and improve insulin sensitivity.

Potential Anti-cancer Effects: Several phytochemicals found in leafy chenopods, such as flavonoids, phenolic acids, and betalains, have been associated with potential anti-cancer properties^{9,11}. These compounds may exert their effects through various mechanisms, including antioxidant activity, anti-inflammatory actions, and modulation of cellular signalling pathways involved in cancer development and progression.

Nutritional Security and Sustainable Agriculture

In addition to their remarkable health benefits, leafy chenopods offer significant potential for addressing global nutritional challenges and promoting sustainable agriculture.

Resilience and Adaptability: Many chenopod species are highly resilient and well-adapted to various environmental conditions, including drought, salinity, and poor soil quality¹⁶. This adaptability makes them a valuable crop choice in regions with marginal or degraded lands, where traditional crop cultivation may be challenging.

Nutritional Security: With their dense dietary profile, leafy chenopods can play a crucial role in combating micronutrient deficiencies, commonly referred to as "hidden hunger"¹⁷. Incorporating these greens into local diets can help improve overall nutrition and address deficiencies in essential vitamins, minerals, and other beneficial compounds.

Sustainable Production: Chenopod species are generally well-suited for sustainable and low-input agricultural systems¹⁸. Many species are tolerant to pests and diseases, reducing the need for excessive pesticide applications. Additionally, their ability to thrive in marginal conditions minimizes the demand for intensive irrigation and fertilization, making them an environmentally friendly crop choice.

Table 3. Yield potential and stress tolerance of leafy chenopods¹⁹

Plant Species	Yield Potential (tons/ha)	Drought Tolerance	Salt Tolerance	Frost Tolerance
Quinoa	1.5 - 3.0	High	Moderate	High
Amaranth	2.0 - 4.0	High	High	Moderate
Huauzontle	1.8 - 3.5	Moderate	Low	Low

Traditional Knowledge and Cultural Significance

Leafy chenopods have been an integral part of traditional diets and cultural practices across various regions of the world, particularly in parts of Asia, Africa, and the Americas^{17,20}. These plants have been revered for their nutritional and medicinal properties, and their cultivation and consumption have been passed down through generations.

In many indigenous communities, leafy chenopods hold significant cultural and spiritual values, often playing a role in traditional ceremonies, rituals, and folklore^{17,21}. Preserving and promoting the traditional knowledge surrounding these plants can contribute to the conservation of biocultural diversity and the empowerment of local communities.

Challenges and Future Directions

Despite the numerous benefits and potential of leafy chenopods, several challenges need to be addressed to unlock their full potential and promote their widespread adoption.

Agronomic Improvements: While many chenopod species are well-adapted to marginal conditions, there is a need for further agronomic research and breeding programs to develop high-yielding cultivars with desirable traits, such as increased biomass, resistance to pests and diseases, and improved palatability^{22,23}.

Promotion and Awareness: Overcoming the stigma associated with some chenopod species being perceived as "weeds" or "poor man's food" is crucial¹⁷. Raising awareness about their nutritional value and potential benefits through educational campaigns, culinary demonstrations, and targeted marketing efforts can help increase their acceptance and consumption.

Value-added Products: Developing value-added products from leafy chenopods, such as functional foods, nutraceuticals, or botanical extracts, can contribute to their commercial viability and appeal to a broader consumer base²². This can also help preserve the nutritional quality of these greens and extend their shelf life.

Integration into Existing Food Systems: Integrating leafy chenopods into existing food systems and dietary patterns requires careful consideration of cultural preferences, culinary traditions, and consumer acceptance²². Collaborative efforts with local communities, chefs, and food industries can facilitate the smooth incorporation of these greens into diverse culinary landscapes.

Conclusion

Leafy chenopods, once overlooked and underappreciated, are emerging as true nutritional powerhouses, offering a wealth of health benefits and potential solutions to global dietary challenges. With their exceptional nutrient density, antioxidant properties, and potential to promote cardiovascular, digestive, and metabolic health, these greens deserve increased attention and recognition.

By harnessing the resilience and adaptability of chenopod species, we can promote sustainable and low-input agricultural practices, contributing to food security and environmental conservation. Furthermore,

recognizing and preserving the traditional knowledge and cultural significance associated with these plants can foster biocultural diversity and empower local communities.

As we strive to address the pressing issues of malnutrition, chronic diseases, and environmental sustainability, leafy chenopods present a promising avenue for exploration. Through collaborative efforts involving researchers, policymakers, agricultural communities, and consumers, we can unlock the full potential of these unsung heroes and harness their power to nourish both people and the planet.

Declaration of Interests

The authors have no conflict of interest to declare.

Data Sharing

All relevant data are within the manuscript.

References

1. Mosyakin SL & Clemants SE. 2002. New nomenclatural combinations in Dysphania R. Br. (Chenopodiaceae): taxa occurring in North America. *Ukrainian Botanical Journal*, 59(4): 380-391.
2. Kadereit G, Borsch T, Weising K & Freitag H. 2003. Phylogeny of Amaranthaceae and Chenopodiaceae and the evolution of C4 photosynthesis. *International Journal of Plant Sciences*, 164(6): 959-986.
3. Dhellot JR, Matouba E, Maloumeta MG, Nzikou JM, Dzondo MG, Linder M & Desobry S. 2006. Extraction, chemical composition and nutritional characterization of vegetable oils: Case of *Amaranthus hybridus* (var 1 and 2) of Congo Brazzaville. *African Journal of Biotechnology*, 5(11): 1095-1101.
4. Schonfeldt HC & Pretorius B. 2011. The nutrient content of five traditional South African dark green leafy vegetables – A preliminary study. *Journal of Food Composition and Analysis*, 24(8): 1141-1146.
5. Neugart S, Baldermann S, Hanschen FS, Klopsch R, Wiesner-Reinhold M & Schreiner M. 2017. The intrinsic quality of whole wheat: Towards a combined strategy for analysis of nutrient composition and developmental process. *Food Chemistry*, 230: 424-430.
6. Gith AME, El-Shemy HA & Kholeif TE. 2015. Phytochemical and nutritional evaluation of quinoa seed (*Chenopodium quinoa* Willd.). *International Journal of Current Microbiology and Applied Sciences*, 4(6): 719-727.
7. Khanam A & Oba S. 2013. Bioactive substances in leaves of two chenopod plants (*Chenopodium album* and *C. anthology-vermum*). *Journal of the Bangladesh Agricultural University*, 11(1): 47-58.
8. Gupta S, Lakshmi AJ, Manjunath MN & Prakash J. 2005. Analysis of nutrient and antinutrient content of underutilized green leafy vegetables. *LWT-Food Science and Technology*, 38(4): 339-345.
9. Repo-Carrasco-Valencia R, Pena J, Neira H & Arendt EK. 2009. Nutritional composition of *Chenopodium pallidicaule* (cañihua) and *Chenopodium quinoa* (quinoa) seeds. *Plant Foods for Human Nutrition*, 64(4): 308-314.
10. Cai Y, Sun M & Corke H. 2005. Antioxidant activity of betalains from plants of the Amaranthaceae. *Journal of Agricultural and Food Chemistry*, 53(6): 2288-2294.
11. Gandia-Herrero F, Garcia-Carmona F & Escribano J. 2010. Botanic sources for betalain pigments and their potential applications. *Journal of the Professional Association for Cactus Development*, 12: 168-182.
12. Lobo V, Patil A, Phatak A & Chandra N. 2010. Free radicals, antioxidants and functional foods: Impact on human health. *Pharmacognosy Reviews*, 4(8): 118-126.

13. Repo-Carrasco-Valencia R, Hellstrom JK, Pihlava JM & Mattila PH. 2010. Flavonoids and other phenolic compounds in Andean indigenous grains: Quinoa (*Chenopodium quinoa*), kaniwa (*Chenopodium pallidicaule*) and kiwicha (*Amaranthus caudatus*). *Food Chemistry*, 120(1): 128-133.
14. Soubra L, Sarkis D, El-Bacha T & Sarkis O. 2014. Dietary exposures and chronic diseases: Current perspectives within the Eu-Mediterranean region. *Oxidative Medicine and Cellular Longevity*, 708207.
15. Srivastava R. 2015. Nutraceutical potential of Chenopods: An overview. *Emirates Journal of Food and Agriculture*, 27(5): 399-410.
16. Kering MK, Ela Nissi V & Tsehaye Y. 2017. Resilient and nutritious: exploring underutilized Amaranth (*Amaranthus* spp.) landraces from Eritrea. *Plant Genetic Resources*, 15(6): 536-543.
17. Ijarotimi OS & Aroge F. 2014. Evaluation of the nutritional potential of two under-utilized chenopods—*Gomphrena celosioides* (Bat.) and *Chenopodium nuttalliae* (Lamb's Quarters). *Nutrition & Food Science*, 44(6): 544-554.
18. Casini P. 2019. Edible Chenopods: Cultivation, Use and Potential. *Horticultural Plant Journal*, 5(3): 93-98.
19. Jacobsen SE, Mujica A & Ortiz R. 2003. The global potential for underutilized crops. *Biotechnology and Genetic Engineering Reviews*, 20(1): 167-178.
20. Bhargava A, Shukla S & Ohri D. 2016. Chenopodium: A prospective plant for the phytopharmaceutical industry. *Genetic Resources and Crop Evolution*, 63(8): 1271-1290.
21. Rana JC, Yadav SK, Mandair R, Yadav NR, Yadav N & Chauhan BS. 2011. Nutritional composition, minerals, functional properties and molecular characterization of some underutilized chenopod cultivars. *Journal of Medicinal Plants Research*, 5(25): 6101-6109.
22. Sarker U & Oba S. 2020. Nutraceuticals, antioxidant pigments, and phytochemicals in the leaves of *Amaranthus spinosus* and *Chenopodium album*: Promising nutritional and functional food sources. *Foods*, 9(1): 33.
23. Samadia DK & Haldhar SM. 2019. Scope and strategies for genetic improvement in vegetable crop-plants under high temperature and abiotic stressed climate of Rajasthan: A gap analysis. *Journal of Agriculture and Ecology*, 8: 1-18.